

(d) *Conditions of use*—(1) *Cattle*—(i) *Dose*. 10 to 15 milligrams per kilogram of body weight, at 24-hour intervals for 3 to 5 consecutive days.

(ii) *Indications for use*. For the treatment of bovine respiratory disease (pneumonia) associated with *Pasteurella haemolytica*, *P. multocida*, and *Haemophilus somnus*.

(iii) *Limitations*. For subcutaneous injection in the neck. Do not inject more than 50 milliliters at each site. Do not slaughter within 11 days of last treatment. Do not use in female dairy cattle 20 months of age or older. Use in this class of cattle may cause residues in milk. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[63 FR 24107, May 1, 1998]

**§ 522.2150 Stanozolol sterile suspension.**

(a) *Specifications*. Each milliliter of sterile suspension contains 50 milligrams of stanozolol.

(b) *Sponsor*. No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) Used as an anabolic steroid treatment in dogs, cats, and horses.

(2) Administered to dogs and cats by deep intramuscular injection in the thigh at weekly intervals, for several weeks. For cats and small breeds of dogs, 25 milligrams. For larger dogs, 50 milligrams.

(3) Administered to horses by deep intramuscular injection in the gluteal region at weekly intervals, for not more than 4 weeks; 25 milligrams per 100 pounds of body weight.

(4) Not for use in horses intended for food.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 46101, Oct. 6, 1975, as amended at 42 FR 36995, July 19, 1977; 55 FR 23076, June 6, 1990]

**§ 522.2200 Sulfachlorpyridazine.**

(a) *Chemical name*. *N*-(6-Chloro-3-pyridazinyl) sulfanilamide.

(b) *Specifications*. Melting point range 190 °C to 191 °C.

(c) *Sponsor*. See No. 053501 in § 510.600(c) of this chapter.

(d) *Related tolerances*. See § 556.630 of this chapter.

(e) *Conditions of use*. It is used for injection into calves as follows:

(1) *Amount*. 30 to 45 milligrams per pound of body weight per day.

(2) *Indications for use*. Treatment of diarrhea caused or complicated by *E. coli* (colibacillosis).

(3) *Limitations*. Administer as the sodium salt of sulfachlorpyridazine intravenously in aqueous solution for 1 to 5 days in divided doses twice daily; treated calves must not be slaughtered for food during treatment or for 5 days after the last treatment.

[40 FR 13858, Mar. 27, 1975, as amended at 50 FR 41489, Oct. 11, 1985]

**§ 522.2220 Sulfadimethoxine injection.**

(a)(1) *Specifications*. Sulfadimethoxine injection containing 400 milligrams per milliliter.

(2) *Sponsor*. (i) See No. 000069 in § 510.600(c) of this chapter for conditions of use as in paragraphs (a)(3)(i) through (a)(3)(iii) of this section.

(ii) See No. 057561 for conditions of use as in paragraph (a)(3) of this section.

(iii) See No. 059130 for use as in paragraph (a)(3)(iii) of this section.

(3) *Conditions of use*. (i) It is used or intended for use in dogs and cats as follows:

(a) For the treatment of respiratory, genitourinary tract, enteric, and soft tissue infections when caused by Streptococci, Staphylococci, Escherichia, Salmonella, Klebsiella, Proteus, or Shigella organisms sensitive to sulfadimethoxine, and in the treatment of canine bacterial enteritis associated with coccidiosis and canine Salmonellosis.

(b) It is administered by intravenous or subcutaneous injection at an initial dose of 55 milligrams per kilogram of body weight followed by 27.5 milligrams per kilogram of body weight every 24 hours.

(c) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

## § 522.2240

(ii) It is used or intended for use in horses as follows:

(a) For the treatment of respiratory disease caused by *Streptococcus equi* (strangles).

(b) It is administered by intravenous injection at an initial dose of 55 milligrams per kilogram of body weight followed by 27.5 milligrams per kilogram of body weight every 24 hours until the patient is asymptomatic for 48 hours.

(c) Not for use in horses intended for food.

(d) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(iii) It is used or intended for use in cattle as follows:

(a) For the treatment of shipping fever complex, bacterial pneumonia, calf diphtheria, and foot-rot.

(b) It is administered by intravenous injection at an initial dose of 25 milligrams per pound of body weight followed by 12.5 milligrams per pound of body weight every 24 hours until the animal is asymptomatic for 48 hours.

(c) Milk taken from animals during treatment and for 60 hours (5 milkings) after the latest treatment must not be used for food. Do not administer within 5 days of slaughter.

(d) Tissue damage may result from perivascular infiltration.

(b) [Reserved]

(c)(1) *Specifications*. Sulfadimethoxine containing 100 milligrams per milliliter.

(2) *Sponsor*. See No. 000010 in § 510.600(c) of this chapter.

(3) *Conditions of use*. (i) It is used or intended for use in the treatment of sulfadimethoxine-susceptible bacterial infections in dogs.

(ii) It is administered by subcutaneous, intramuscular, or intravenous injection at an initial dose of 25 milligrams per pound of body weight followed by 12.5 milligrams per pound of body weight every 24 hours thereafter. Continue treatment until the animal is free from symptoms for 48 hours.

(iii) For use by or on the order of a licensed veterinarian.

## 21 CFR Ch. I (4–1–05 Edition)

(d) *Related tolerances*. See § 556.640 of this chapter.

[40 FR 13858, Mar. 27, 1975, as amended at 40 FR 34112, Aug. 14, 1975; 40 FR 42007, Sept. 10, 1975; 50 FR 254, Jan. 3, 1985; 53 FR 40728, Oct. 18, 1988; 54 FR 30205, July 19, 1989; 58 FR 38972, July 21, 1993; 59 FR 56000, Nov. 10, 1994; 61 FR 4875, Feb. 9, 1996; 62 FR 23128, Apr. 29, 1997; 62 FR 35076, June 30, 1997]

### § 522.2240 Sulfathoxypyridazine.

(a) *Chemical name*. N<sup>1</sup>-(6-Ethoxy-3-pyridazinyl) sulfanilamide.

(b) *Specifications*. Melting point range of 180 °C to 186 °C.

(c) *Sponsor*. See No. 053501 in § 510.600(c) of this chapter.

(d) *Related tolerances*. See § 556.650 of this chapter.

(e) *Conditions of use*. It is used for injection into cattle as follows:

(1) *Amount*. 2.5 grams per 100 pounds of body weight per day.

(2) *Indications for use*. Treatment of respiratory infection (pneumonia, shipping fever), foot rot, calf scours; as adjunctive therapy in septicemia accompanying mastitis and metritis.

(3) *Limitations*. Administer intravenously for not more than 4 days; or first treatment may be followed by 3 days of treatment with sulfathoxypyridazine in drinking water, feed, or tablet in accordance with § 558.579(e) or §§ 520.2240a(e) and 520.2240b(e) of this chapter; as sodium sulfathoxypyridazine; do not treat within 16 days of slaughter; as sole source of sulfonamide; milk that has been taken from animals during treatment and for 72 hours (6 milkings) after the latest treatment must not be used for food; for use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 41 FR 11011, Mar. 15, 1976; 67 FR 78355, Dec. 24, 2002]

### § 522.2260 Sulfamethazine injectable solution.

(a) *Specifications*. Each milliliter of sterile aqueous solution contains 250 milligrams of sulfamethazine sodium.

(b) *Sponsor*. See No. 053501 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.670 of this chapter.

(d) *NAS/NRC status*. The conditions of use are NAS/NRC reviewed and found